

Organism's responses to a long-term inhalation of silica-containing submicron particles (predominantly, nanoscale) of an industrial aerosol

S.N. Solovyeva¹, B.A. Katsnelson¹, M.P. Sutunkova¹, L.I. Privalova¹,
V.B. Gurvich¹, I.A. Minigalieva¹, T.V. Slyshkina¹, I.E. Valamina²,
V.Ya. Shur³, I.V. Zubarev³, D.K. Kuznetsov³

¹*Yekaterinburg Medical Research Center for Prophylaxis and Health Protection in Industrial Workers, 620014, Yekaterinburg, Russia
solovyevasn@ymrc.ru*

²*Central Research Laboratory of Ural State Medical University, 620028, Yekaterinburg, Russia*

³*School of Natural Sciences and Mathematics, Ural Federal University, 620000, Yekaterinburg, Russia*

Outbred female white rats were exposed for 3 or 6 months, 5 times a week, 4 h a day to an aerosol containing 72% amorphous silica predominantly submicron particles (nanoscale ones included) at an exposure concentration of 2.6 ± 0.6 or 10.6 ± 2.1 mg/m³.

The silica (mostly amorphous) containing submicron particles with a prevailing proportion of those in the nanoscale range induce, when instilled intratracheally, a pulmonary cell response comparable with that to highly cytotoxic and fibrogenic standard quartz powder DQ12. Nevertheless, in long-term inhalation experiment at realistic concentrations, they proved to be of very low systemic toxicity and negligible pulmonary fibrogenicity.

This paradox may be explained by low SiO₂ retention in lungs and other organs due to a relatively high solubility of these nanoparticles in relevant biological and model milieus.